




Alaska Fisheries Management and Climate Change




Gretchen A. Harrington and Julie A. Scheurer, NMFS Alaska Region

Climate Change

- Rising mean global temperatures
- Faster and larger changes are happening in the northern latitudes
- Greatest mean surface temperature anomalies (darkest red) are occurring in the Arctic, as shown in figure below



http://hydrogencommerce.com/image/polar_bears480.jpg



An Ice-Free Alaska

- Opens entire 47,000 miles of coastline
- Increases fishable marine waters off Alaska from approximately 360,000 to 610,000 square miles

What do Scientists Think This Means for Alaska?

Ecological Surprise!

- Restructuring of Alaska marine ecosystems
- Changes in physical and chemical properties of oceans (changing currents, ocean acidification...)
- Shifts in species abundance and distribution
- Expanding ranges for warm water species (arrowtooth flounder, mackerel, spiny dogfish...)
- Contracting ranges for cold water and ice-dependent species (marine mammals, seabirds, salmon, crab...)
- Increases in coastal erosion

Disappearing Sea Ice



Maps show the shrinking winter and summer ice edges.

Preparing for the Future: Management in a Changing Marine Environment



<http://www.aaas.org/news/releases/2006/>

Changing Human Actions


- Changing fishing practices:
 - Catch different species
 - Open new, previously unfished, areas
 - Fish further from ports/processors
 - Use larger vessels with capacity to travel greater distances
- Changing subsistence harvests by local Native people
- Increasing potential for coastal development
- Opening trans-Arctic shipping routes

Bering Sea Fishery Management

- The North Pacific Fishery Management Council has recommended amendments to the Fishery Management Plan to address the northward expansion of fisheries and uncertainty in changing environment:
 - Freeze footprint of existing bottom trawl fisheries
 - Close areas of important habitat
 - Close areas important for subsistence uses
 - Establish the Northern Bering Sea Research Area to investigate the effects of bottom trawling on benthic habitat

Arctic Fishery Management

- Marine waters currently inaccessible and marine resources poorly described
- First step - Prohibit commercial fisheries until scientific knowledge supports fishery development and management
- Second step - Develop a new Arctic Fishery Management Plan to manage marine resources in the Chukchi and Beaufort Seas considering:
 - Opportunities
 - Act precautionary
 - Develop Fishery Ecosystem Plan
 - Use adaptive management
 - Involve Native communities
 - Challenges
 - Limited abundance and distribution data
 - High uncertainty of long-term effects
 - Risk of human actions compounding the effects
 - Potential for declines in marine mammals and seabirds



CELESTIAL FOCUS/ONYX

Acknowledgements: Thanks to Dr. Gordon Kruse (University of Alaska Fairbanks), Mike Litzow (NMFS Alaska Fishery Science Center), Bill Wilson (North Pacific Fishery Management Council), and Steve Lewis (NMFS Alaska Region) for the information used in this poster.

<http://www.cnn.com/CNN/Programs/presents/shows/melting.point/images>

U.S. mission heads north to map Arctic continental shelf

CROWDED: Polar waters draw Denmark, Russia, Canada also.

By DOUG ESSER
The Associated Press

SEATTLE -- A U.S. Coast Guard icebreaker is headed to the Arctic to map the seafloor off Alaska, as Russia, Denmark and Canada assert their claims in the polar region, which has potential oil and gas reserves.

The lead scientist on the expedition scoffs at the political implications.

"We're basically just doing science," said Larry Mayer, director of the Center for Coastal and Ocean Mapping at the University of New Hampshire.

"There's no flag-dropping on this trip," he said Thursday in an interview from Durham, N.H.

Mayer will meet the Seattle-based icebreaker Healy at Barrow and head about 500 miles north with a team of about 20 scientists to map an area known as the Chukchi Cap.

The Healy left Puget Sound last Monday and should be in Barrow around Friday, said Russ Tippets, a spokesman at the Coast Guard Pacific area office in San Francisco.

"We've had this trip planned for months and it has nothing to do with the Russians planting their flag," he said Thursday in an interview.

Russia has made no official comment on the Healy's trip, but Russian media have suggested it's a response to a Russian submarine dropping that nation's flag Aug. 2 on the floor of the Arctic Ocean under the North Pole.

Russian media say the Healy's mission signals the United States, along with Canada, is actively joining the competition for resources in the Arctic.

Melting ice could open water for drilling or create the long-sought Northwest Passage for shipping.

Canadian Prime Minister Stephen Harper went to the Arctic last week to reassert sovereignty after the Russian flag incident. Canada announced plans in July to build up to eight icebreakers to patrol Arctic waters.

Meanwhile, Danish scientists head for the ice pack Sunday on the Swedish icebreaker Oden seeking evidence that the 1,240-mile underwater Lomonosov Ridge is attached to Greenland, making it a geological extension of Denmark's territory.

The Russians are gathering evidence to support a claim Russia owns the Lomonosov Ridge.

The purpose of the mapping work aboard the Healy is to determine the extent of the continental shelf north of Alaska, Mayer said.

"In that area the country would have rights over the resources of the sea floor and subsurface that would include drilling for oil and gas," he said.

It's not a claim, he said, but a process of registering boundary information with the United Nation's Commission on the Limits of the Continental Shelf.

The mapping is conducted with an echo sounder, something that's difficult to hear when a ship is crunching through the ice, Mayer said. This is the third such mapping trip. The others were in 2003 and 2004.

There will be about 20 scientists on board the Healy with the crew, including representatives from the University of New Hampshire, the University of Texas, the University of Alaska, the National Ice Center in Suitland, Md., the Scripps Institution, the National Oceanic and Atmospheric Administration, and the State Department, he said.

The State Department says its official is a geographer and member of an extended continental shelf task force.

"While a significant technological achievement, the planting of the Russian flag on the seabed of the North Pole has no legal effect and did not prompt the participation of the State Department expert in the Healy cruise," said spokeswoman Nicole Thompson.

The Healy's mission will last a couple of months. It's due back in Seattle in early October, said spokesman Stephen Elliott.

The seven-year-old Healy is the nation's newest icebreaker. It's 420 feet long and is capable of breaking ice 8 feet thick.

Anchorage Daily News
August 13, 2007

Read the sunspots

The mud at the bottom of B.C. fjords reveals that solar output drives climate change - and that we should prepare now for dangerous global cooling

R. TIMOTHY PATTERSON

Financial Post

Wednesday, June 20, 2007


Politicians and environmentalists these days convey the impression that climate-change research is an exceptionally dull field with little left to discover. We are assured by everyone from David Suzuki to Al Gore to Prime Minister Stephen Harper that "the science is settled." At the recent G8 summit, German Chancellor Angela Merkel even attempted to convince world leaders to play God by restricting carbon-dioxide emissions to a level that would magically limit the rise in world temperatures to 2C.

The fact that science is many years away from properly understanding global climate doesn't seem to bother our leaders at all. Inviting testimony only from those who don't question political orthodoxy on the issue, parliamentarians are charging ahead with the impossible and expensive goal of "stopping global climate change." Liberal MP Ralph Goodale's June 11 House of Commons assertion that Parliament should have "a real good discussion about the potential for carbon capture and sequestration in dealing with carbon dioxide, which has tremendous potential for improving the climate, not only here in Canada but around the world," would be humorous were he, and even the current government, not deadly serious about devoting vast resources to this hopeless crusade.

Climate stability has never been a feature of planet Earth. The only constant about climate is change; it changes continually and, at times, quite rapidly. Many times in the past, temperatures were far higher than today, and occasionally, temperatures were colder. As recently as 6,000 years ago, it was about 3C warmer than now. Ten thousand years ago, while the world was coming out of the thousand-year-long "Younger Dryas" cold episode, temperatures rose as much as 6C in a decade -- 100 times faster than the past century's 0.6C warming that has so upset environmentalists.

Climate-change research is now literally exploding with new findings. Since the 1997 Kyoto Protocol, the field has had more research than in all previous years combined and the discoveries are completely shattering the myths. For example, I and the first-class scientists I work with are consistently finding excellent correlations between the regular fluctuations in the brightness of the sun and earthly climate. This is not surprising. The sun and the stars are the ultimate source of all energy on the planet.

My interest in the current climate-change debate was triggered in 1998, when I was funded by a Natural Sciences and Engineering Research Council strategic project grant to determine if there were regular cycles in West Coast fish productivity. As a result of wide swings in the populations of anchovies, herring and other commercially important West Coast fish stock, fisheries managers were having a very difficult time establishing appropriate fishing quotas. One season there would be abundant stock and broad harvesting would be acceptable; the



CREDIT: Andrew Barr, National Post
(See hardcopy for Chart/Graph)

very next year the fisheries would collapse. No one really knew why or how to predict the future health of this crucially important resource.

Although climate was suspected to play a significant role in marine productivity, only since the beginning of the 20th century have accurate fishing and temperature records been kept in this region of the northeast Pacific. We needed indicators of fish productivity over thousands of years to see whether there were recurring cycles in populations and what phenomena may be driving the changes.

My research team began to collect and analyze core samples from the bottom of deep Western Canadian fjords. The regions in which we chose to conduct our research, Effingham Inlet on the West Coast of Vancouver Island, and in 2001, sounds in the Belize-Seymour Inlet complex on the mainland coast of British Columbia, were perfect for this sort of work. The topography of these fjords is such that they contain deep basins that are subject to little water transfer from the open ocean and so water near the bottom is relatively stagnant and very low in oxygen content. As a consequence, the floors of these basins are mostly lifeless and sediment layers build up year after year, undisturbed over millennia.

Using various coring technologies, we have been able to collect more than 5,000 years' worth of mud in these basins, with the oldest layers coming from a depth of about 11 metres below the fjord floor. Clearly visible in our mud cores are annual changes that record the different seasons: corresponding to the cool, rainy winter seasons, we see dark layers composed mostly of dirt washed into the fjord from the land; in the warm summer months we see abundant fossilized fish scales and diatoms (the most common form of phytoplankton, or single-celled ocean plants) that have fallen to the fjord floor from nutrient-rich surface waters. In years when warm summers dominated climate in the region, we clearly see far thicker layers of diatoms and fish scales than we do in cooler years. Ours is one of the highest-quality climate records available anywhere today and in it we see obvious confirmation that natural climate change can be dramatic. For example, in the middle of a 62-year slice of the record at about 4,400 years ago, there was a shift in climate in only a couple of seasons from warm, dry and sunny conditions to one that was mostly cold and rainy for several decades.

Using computers to conduct what is referred to as a "time series analysis" on the colouration and thickness of the annual layers, we have discovered repeated cycles in marine productivity in this, a region larger than Europe. Specifically, we find a very strong and consistent 11-year cycle throughout the whole record in the sediments and diatom remains. This correlates closely to the well-known 11-year "Schwabe" sunspot cycle, during which the output of the sun varies by about 0.1%. Sunspots, violent storms on the surface of the sun, have the effect of increasing solar output, so, by counting the spots visible on the surface of our star, we have an indirect measure of its varying brightness. Such records have been kept for many centuries and match very well with the changes in marine productivity we are observing.

In the sediment, diatom and fish-scale records, we also see longer period cycles, all correlating closely with other well-known regular solar variations. In particular, we see marine productivity cycles that match well with the sun's 75-90-year "Gleissberg Cycle," the 200-500-year "Suess Cycle" and the 1,100-1,500-year "Bond Cycle." The strength of these cycles is seen to vary over time, fading in and out over the millennia. The variation in the sun's brightness over these longer cycles may be many times greater in magnitude than that measured over the short Schwabe cycle and so are seen to impact marine productivity even more significantly.

Our finding of a direct correlation between variations in the brightness of the sun and earthly climate indicators (called "proxies") is not unique. Hundreds of other studies, using proxies from tree rings in Russia's Kola Peninsula to water levels of the Nile, show exactly the same thing: The sun appears to drive climate change.

However, there was a problem. Despite this clear and repeated correlation, the measured variations in incoming solar energy were, on their own, not sufficient to cause the climate changes we have observed in our proxies. In addition, even though the sun is brighter now than at any time in the past 8,000 years, the increase in direct solar input is not calculated to be sufficient to cause the past century's modest warming on its own. There had to be an amplifier of some sort for the sun to be a primary driver of climate change.

Indeed, that is precisely what has been discovered. In a series of groundbreaking scientific papers starting in 2002, Veizer, Shaviv, Carslaw, and most recently Svensmark et al., have collectively demonstrated that as the output of the sun varies, and with it, our star's protective solar wind, varying amounts of galactic cosmic rays from deep space are able to enter our solar system and penetrate the Earth's atmosphere. These cosmic rays enhance cloud formation which, overall, has a cooling effect on the planet. When the sun's energy output is greater, not only does the Earth warm slightly due to direct solar heating, but the stronger solar wind generated during these "high sun" periods blocks many of the cosmic rays from entering our atmosphere. Cloud cover decreases and the Earth warms still more.

The opposite occurs when the sun is less bright. More cosmic rays are able to get through to Earth's atmosphere, more clouds form, and the planet cools more than would otherwise be the case due to direct solar effects alone. This is precisely what happened from the middle of the 17th century into the early 18th century, when the solar energy input to our atmosphere, as indicated by the number of sunspots, was at a minimum and the planet was stuck in the Little Ice Age. These new findings suggest that changes in the output of the sun caused the most recent climate change. By comparison, CO₂ variations show little correlation with our planet's climate on long, medium and even short time scales.

In some fields the science is indeed "settled." For example, plate tectonics, once highly controversial, is now so well-established that we rarely see papers on the subject at all. But the science of global climate change is still in its infancy, with many thousands of papers published every year. In a 2003 poll conducted by German environmental researchers Dennis Bray and Hans von Storch, two-thirds of more than 530 climate scientists from 27 countries surveyed did not believe that "the current state of scientific knowledge is developed well enough to allow for a reasonable assessment of the effects of greenhouse gases." About half of those polled stated that the science of climate change was not sufficiently settled to pass the issue over to policymakers at all.

Solar scientists predict that, by 2020, the sun will be starting into its weakest Schwabe solar cycle of the past two centuries, likely leading to unusually cool conditions on Earth. Beginning to plan for adaptation to such a cool period, one which may continue well beyond one 11-year cycle, as did the Little Ice Age, should be a priority for governments. It is global cooling, not warming, that is the major climate threat to the world, especially Canada. As a country at the northern limit to agriculture in the world, it would take very little cooling to destroy much of our food crops, while a warming would only require that we adopt farming techniques practiced to the south of us.

Meantime, we need to continue research into this, the most complex field of science ever tackled, and immediately halt wasted expenditures on the King Canute-like task of "stopping climate change."

R. Timothy Patterson is professor and director of the Ottawa-Carleton Geoscience Centre, Department of Earth Sciences, Carleton University.